

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property  
Organization  
International Bureau



(43) International Publication Date  
21 October 2004 (21.10.2004)

PCT

(10) International Publication Number  
**WO 2004/089369 A3**

- (51) International Patent Classification<sup>7</sup>: **A61K 31/436**,  
A61P 25/16, 25/28, 43/00, G01N 33/50, 33/68
- (21) International Application Number:  
PCT/GB2004/000690
- (22) International Filing Date: 24 February 2004 (24.02.2004)
- (25) Filing Language: English
- (26) Publication Language: English
- (30) Priority Data:  
60/462,269 11 April 2003 (11.04.2003) US
- (71) Applicant (for all designated States except US): **CAMBRIDGE UNIVERSITY TECHNICAL SERVICES LIMITED** [GB/GB]; The Old Schools, Trinity Lane, Cambridge, Cambridgeshire CB2 1TS (GB).
- (72) Inventors; and
- (75) Inventors/Applicants (for US only): **RUBINSZTEIN, David** [GB/GB]; Cambridge Institute For Medical Research, Wellcome/MRC Building, Addenbrookes Hospital, Hill Road, Cambridge, Cambridgeshire CB2 2XY (GB). **RAVIKUMAR, Brinda** [IN/GB]; Cambridge Institute For Medical Research, Wellcome/MRC Building, Addenbrookes Hospital, Hill Road, Cambridge, Cambridgeshire CB2 2XY (GB). **WEBB, Julie** [GB/GB]; Cambridge Institute For Medical Research, Wellcome/MRC Building, Addenbrookes Hospital, Hill Road, Cambridge, Cambridgeshire CB2 2XY (GB).
- (81) Designated States (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NA, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW.
- (84) Designated States (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PT, RO, SE, SI, SK, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).
- Published:  
— with international search report  
— with amended claims
- (88) Date of publication of the international search report:  
7 April 2005
- Date of publication of the amended claims: 23 June 2005
- For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

(54) Title: METHODS AND MEANS FOR TREATING PROTEIN CONFORMATIONAL DISORDERS

(57) Abstract: This invention relates to the recognition that autophagy plays a key role in the clearance of the intracellular protein aggregates which characterise Protein Conformational Disorders, such as Huntington's disease and Parkinson's disease. Methods and uses of autophagy inducing agents, such as rapamycin macrolides, in the treatment of Protein Conformational Disorders, are described herein.

WO 2004/089369 A3

## AMENDED CLAIMS

[Received by the International Bureau on 08 July 2004 (08.07.2004):  
original claims 38 and 40 amended; (2 pages)]

30. Use according any one of claims 17 to 23 wherein the disorder is a prion disorder.
- 5 31. Use according to claim 30 wherein the prion disorder is CJD.
32. A method of identifying an agent useful in the treatment of a protein conformational disorder comprising;
- 10 contacting a mammalian cell with a test compound; and, determining the autophagy activity of said cell, an increase in autophagy activity in the presence of said compound being indicative that the compound is a candidate agent for use in the treatment of a protein conformational
- 15 disorder.
33. A method according to claim 32 wherein the cell comprises a heterologous nucleic acid encoding an aggregation-prone polypeptide.
- 20
34. A method according to claim 33 wherein said heterologous nucleic acid is operably linked to an inducible promoter.
35. A method according to claim 33 or claim 34 comprising
- 25 expressing said nucleic acid and stopping said expression, prior to contacting the mammalian cell with the test compound.
36. A method according to any one of claims 32 to 35 comprising modifying the compound to optimise the
- 30 pharmaceutical properties thereof
37. A method according to any one of claims 32 to 36 comprising formulating the test compound into a pharmaceutical composition.
- 35
38. A method of producing an agent for the treatment of a protein conformational disorder comprising;

modifying rapamycin to produce a rapamycin derivative; and;  
determining the autophagy inducing activity of said  
derivative.

5 39. A method according to claim 38 comprising determining the  
ability of said derivative to inhibit mTOR.

40. A method according to claim 38 or claim 39 comprising  
determining the ability of said derivative to enhance the  
10 clearance of cytoplasmic protein aggregates.